



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A01N 3/02, A47G 7/02, B65D 85/50, A01G 5/06	A1	(11) International Publication Number: WO 99/55154 (43) International Publication Date: 4 November 1999 (04.11.99)
(21) International Application Number: PCT/NL99/00239 (22) International Filing Date: 26 April 1999 (26.04.99) (30) Priority Data: 1008986 24 April 1998 (24.04.98) NL (71)(72) Applicants and Inventors: VONK, Hans [NL/NL]; Lange Schilk 5, NL-2461 LC Ter Aar (NL). STAR, Tinus, Willem, Hendrik [NL/NL]; Het Cappelrijeland 9, NL-2231 ZD Rijsburg (NL). (74) Agent: ALTENBURG, Bernardus, Stephanus, Franciscus; Octrooibureau Los en Stigter B.V., Weteringschans 96, NL-1017 XS Amsterdam (NL).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i> <i>In English translation (filed in Dutch).</i>
(54) Title: METHOD OF PRESERVING CUT FLOWERS, GEL ON WATER BASIS AND A HOLDER		
(57) Abstract The invention relates to a method of preserving cut flowers, wherein the cut flowers are placed with their stems into a holder containing a gel on water basis. According to the invention gel on polyacrylamide basis is used in which the size of the dry granules is 0.2 mm or smaller. The invention also relates to such a gel and a holder suitable for the preservation of cut flowers in the gel.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

Method of preserving cut flowers, gel on water basis and a holder

The present invention relates to a method of preserving cut flowers, wherein the cut flowers are placed with their stems into a holder, which holder is provided with a bottom and contains an aqueous gel as medium, and
5 the gel is prepared by allowing granules of a water absorbent cross-linked polymer to swell.

It is known that to preserve cut flowers they generally have to stand substantially continuously in water. This also applies during transport, especially for the
10 most delicate cut flowers such as lathyrus. During the transport of holders filled with water the water may splash out, or the holders may fall over so that the cut flowers stand dry. At the same time, bacteria and moulds and consequently diseases as, for example, botrytis may
15 spread through the water. A further disadvantage is that the cut flowers have to be transported vertically, which means that more space is required. This in turn raises the transport costs. It makes transport of cut flowers over large distances very expensive.

20 To solve this problem the prior art proposes the use of a gel (see, for example, SE-8403211). The known gels exhibit various disadvantages. Some gels themselves serve as medium for bacteria, thereby increasing the problem of spreading of disease. The advantage of gels based
25 on swollen granules is that if they inadvertently become distributed over plant parts, only a smaller part will become infected compared with a gel based on a dissolved polymer. From practice and from literature the use of gels based on granules with a size of 0.5 mm is already known.
30 The interstitial spaces between the swollen granules contain liquid water. This water is still able to spread diseases, albeit to a limited degree. For this reason it is a known practice in the prior art to dissolve a polymer in said interstitial water and to make the water into gel.
35 Such a gel is relatively complex (different gel forming

components) and has been shown in practice to be difficult to handle, for example, when transferring the gel into a holder. If a gel is used that has been prepared by adding to the granules only so much water as the granules are
5 able to absorb, a cut flower that has been inserted into the gel will be unable to take up sufficient water.

It is the aim of the present invention to eliminate the above-mentioned disadvantages to a large extent.

To this end the method according to the present
10 invention is characterized in that the stems are inserted into a gel on polyacrylamide basis, in which the size of the dry granules is 0.2 mm or smaller.

Surprisingly it has been shown that a gel based on a polyacrylamide-comprising polymer, in which the granules
15 in dry condition have a size of 0.2 mm or smaller, is not only easier to handle (i.e. easier to pump), but also that there is no longer any need for a polymer in solution. Moreover, it appears that sufficient water is released for the preservation of the cut flowers. Without wishing to be
20 bound to any theory, it is believed that due to their larger surface area smaller granules are better able to release water to be taken up by the cut flowers. The gel used in the method is liquid, that is to say it is able to flow. The lower-limit size of the granules does not appear
25 to be typical, but preferably it will in general be larger than the diameter of the capillaries of the respective cut flower. Although a soluble polymer may be used, it is not required and by refraining from doing so, a simple gel-forming composition may be provided.

30 A gel suitable for application in the method according to the present invention contains sufficient gel-forming polymer to increase the viscosity of the aqueous medium such that it will not easily splash or run out of the holder. The upper limit of the concentration of
35 gel-forming polymer in a gel according to the invention is determined by the ability of the gel to release water to the cut flower. With the method according to the present invention there is no objection to the gel exhibiting some syneresis, that is to say that it temporally loses water.

Depending on the desired flow behaviour and the storage life of the cut flowers, a person knowledgeable about gels is able in a simple manner to determine a suitable concentration of gel-forming polymer. If a higher
5 concentration unacceptably shortens the storage life, the concentration of gel-forming polymer has to be reduced. If the aqueous medium is too thin, the concentration of gel-forming polymer has to be increased. Reference in the present application to a granule size of less than 0.2 mm is
10 understood to mean a size distribution of granules, in which more than 80% by weight, and preferably at least 90% by weight of the granules have a size of 0.2 mm or smaller.

A typical suitable concentration of gel-forming
15 polymer is 3 to 25 g, preferably between 5 and 20 g per litre of the final gel.

According to a preferred embodiment, the gel comprises 6 to 14 g gel-forming polymer per litre of the final gel.

20 According to a favourable embodiment the aqueous gel further comprises at least one compound chosen from the group comprised of a disinfectant, a surfactant, nutrients and a colouring agent.

The first three promote the storage life of the cut
25 flowers. A colouring agent may, for example, be used for the identification of a gel with a particular composition, for example, to indicate for which kind of cut flowers the gel is suitable.

According to a very favourable embodiment the aqueous
30 gel comprises polyacrylamide and has a pH between 2.5 and 7, preferably between 4 and 6.5.

Such gels have been proven to be very effective for the preservation of cut flowers.

The method according to the present invention is in
35 particular also suitable for the preservation of cut flowers during transport. The cut flowers are preferably transported in a substantially horizontal position.

As a result it has become possible to also transport cut flowers that are sensitive to dryness using up a smaller volume. This means that air transport is more often economically feasible, thereby allowing the supply to markets that could previously not be catered for because of the short life that would remain for the flowers once they have reached the consumer. For nearer markets applies that the consumer is able to enjoy the flowers longer. The term "substantially horizontally" is understood to mean an angle smaller than approximately 25°.

The present invention also relates to a gel on water basis provided with nutrients for cut flowers, which gel has been prepared by allowing granules of a water-absorbent cross-linked polymer on polyacrylamide basis to swell, which granules in dry state have a size of 0.2 mm or smaller.

Finally, the present invention relates to a holder for the preservation of cut flowers, which holder comprises a bottom and an opening for the insertion of the stem of the cut flower and is provided with a gel according to the invention, which holder is further provided with means for keeping the end of a stem of the cut flower at a distance from the bottom.

Especially when the stem of the cut flower has been cut straight across, there is a risk when using the gel according to the present invention that due to contact with the holder the cut-off end does not make sufficient contact with the gel granules, so that the cut flower does not receive an adequate amount of water. This is particularly a problem with the gel according to the present invention because of the smaller diameter of the water-conducting channels between the swollen gel granules (capillary forces). The provision of means for distancing ensures that sufficient gel granules make contact with the cut end of the flower.

According to a preferred embodiment the holder is provided with sealing means chosen from the group comprising i) a cap sealing the opening by means of clamps; ii) a

pierceable membrane; and iii) a removable membrane provided with means for removing the membrane by pulling it off.

Such a holder can be prefabricated and can simply
5 be prepared for use for the preservation of a cut flower by a flower grower, florist or consumer .

The invention will be explained with reference to the following exemplary embodiments and with reference to the drawings in which

10 Fig. 1 represents a longitudinal section of a holder according to the invention; and

Fig. 2 represents a cross-section of the holder of Fig. 1 along line II-II.

15 1) Preparation of a gel for the preservation of cut flowers.

76 g polyacrylamide having a granule size < 0.2 mm, Stockosorb 400 F (Chemische Fabrik Stockhausen GmbH, Krefeld, Germany) which is 0.1% cross-linked, is added to 10 l of a solution comprised of 5 g/l Flower food (Laboratorium van der Sprong B.V., Roelofarendsveen, The Netherlands) for lathyrus, or 10 g/l for carnations. Among other things, Flower food comprises a disinfectant, a surfactant, and nutrients. For the person skilled in the art it will be obvious that depending on the crop to be pre-
25 served, other compositions that are known as such, may be used. The mixture is blended until a homogenous gel is obtained. The pH of the gel is approximately 5.

2) The gel's suitability for lathyrus.

The suitability of the gel was tested by placing
30 lathyrus cut flowers in the gel as prepared above. As control, an aqueous medium was used having the same composition but without the gel-forming agent.

The presence of the gel-forming agent appeared to have no adverse effect on the storage life.

35 3) The gel's suitability for carnations.

The suitability of the gel according to the invention was tested by placing carnations into the gel prepared as in 1). As control, an aqueous medium was used

having the same composition but without the gel-forming agent. Without the gel-forming agent the storage life was 12.8 days and with the gel-forming agent 12.4 days.

4) The effect of gel granule size.

5 In the manner described for 1) for lathyrus, gels were made on the basis of

- Stockosorb 400F (0-200 micron)
- Stockosorb 400F (200-800 micron)
- Stockosorb 400RD (100-800 micron)
- 10 - Stockosorb 410K (800-3000 micron)

As control flowers from the same batch were used without being placed into a gel. All flowers were stored in a horizontal position for 48 hours at 16°C. After completion of this 'simulation of transport' all the flowers, those in
15 gel as well as those of the control group, were wilted and discoloured. However, this did not apply to the flowers that were stored in the gel comprising Stockosorb 400F. These were still completely turgid and had their original colour. Subsequently, the storage life of all the bunches
20 of flowers was examined by placing the individual bunches into a vase with clean water. The storage life of the control group was shown to be the same as that of the flowers tested on gels with larger granules. The flowers that were stored on Stockosorb 400F had a longer storage life. As
25 additional controls use is made of gels based on Terawet™ (polyacrylate/polyacrylamide copolymers; granule size between 70 and 2000 micron; Terawet Corp., San Diego, USA) and Broadleaf™ P4 (polyacrylamide gel; precise granule size unknown but relatively large. Greenacres, Denham,
30 UK). These also did not show the favourable results that were seen when using a gel according to the invention.

As holder, a deformable holder may be used, whose end opposite the bottom is pressed against the stems with the aid of fastening means. In this manner the gel is con-
35 tained by the holder, while due to the high viscosity the gel is not, or only with difficulty able to pass through the narrow gaps between the stems. This is an effective way of achieving that the bottom ends of the stems remain in contact with an aqueous medium while the likelihood of

spreading diseases is greatly reduced. Suitable fastening means are, for example, string, a rectangular strip of plastic reinforced with wire, and according to a favourable embodiment, an elastic band or tape. The holder is
5 made of a gel- or water-impermeable material, for example a plastic such as polyethylene. The deformable holder is, for example, a plastic bag.

The holder may also be a rigid holder, such as a holder for one cut flower. These are known to the public
10 to be used mostly for orchids, but may also be used for other cut flowers such as roses and gerberas. In order to ensure supply of water to the cut flower when the gel according to the present invention is used, and in particular in the case of the stem being cut straight-across,
15 the holder has to be provided with means for keeping the end of the stem off the wall and in particular off the bottom of the holder. There are various ways of achieving this. One could consider elements projecting from the wall of the holder, bending the stem in an undulatory manner.
20 This causes the holder to be held to the stem by means of friction. The holder may also be provided with projections on which the end of the stem rests, which projections will to a large extent leave the end of the stem free.

A suitable embodiment is represented in Fig. 1,
25 showing a longitudinal section of a holder 1, with a cavity 2 for the accommodation of gel. The bottom 3 is provided with protuberances 4 leaving open a channel 5. The width of the channel 5 is greater than the size of the swollen gel granules and smaller than the diameter of the
30 stem of the cut flower for which the holder has been designed. In this way a sufficient supply of water to the cut flower is ensured. If the stem happens to come between the wall of the holder 1 and a projection 4 the supply of water is also ensured.

35 Fig. 2 shows a cross-section of the holder 1 wherein the embodiment represented is provided with two channels 5 and four protuberances 4.

For use the holder is conveniently provided with a seal, such as a cap. The cap may be a cap that clamps around the open end of the holder 1, or a cap that clamps against the inside of the opening. In the latter case the cap is conveniently provided with a lip part to facilitate removal of the cap. The holder may also be sealed by means of a membrane, such as plastic foil that is glued to the end. The membrane may be provided with non-perforated grooves, which make it easier to prick a stem through the locally weakened membrane. Alternatively, the membrane may also be removed before use, in which case the membrane is suitably provided with a lip part.

CLAIMS

1. A method of preserving cut flowers, wherein the cut flowers are placed with their stems into a holder, which holder is provided with a bottom and contains an aqueous gel as medium, and the gel is prepared by allowing
5 granules of a water absorbent cross-linked polymer to swell, **characterized** in that the stems are inserted into a gel on polyacrylamide basis, in which the size of the dry granules is 0.2 mm or smaller.
2. A method according to claim 1, **characterized** in
10 that the gel comprises 5 to 20 g gel-forming polymer per litre of the final gel.
3. A method according to claim 2, **characterized** in that the gel comprises 6 to 14 g gel-forming polymer per litre of the final gel.
- 15 4. A method according to one of the preceding claims, **characterized** in that the aqueous gel further comprises at least one compound chosen from the group comprised of a disinfectant, a surfactant, nutrients and a colouring agent.
- 20 5. A method according to one of the preceding claims, **characterized** in that the aqueous gel has a pH between 4.5 and 7.
6. A method for the preservation of cut flowers transport according to one of the preceding claims, **char-**
25 **acterized** in that the cut flowers are transported in a substantially horizontal position.
7. A method according to one of the preceding claims, **characterized** in that a holder is used, which holder is further provided with means for keeping the end
30 of the stem of a cut flower at a distance from the bottom.
8. A gel on water basis provided with nutrients for cut flowers, which gel has been prepared by allowing granules of a water-absorbent cross-linked polymer on polyacrylamide basis to swell, which granules in dry state
35 have a size of 0.2 mm or smaller.

9. A holder for the preservation of cut flowers, which holder comprises a bottom and an opening for the insertion of the stem of the cut flower and is provided with a gel according to claim 8, which holder is further
5 provided with means for keeping the end of a stem of the cut flower at a distance from the bottom.

10. A holder according to claim 9, which holder is provided with sealing means chosen from the group comprising i) a cap sealing the opening by means of clamps; ii) a
10 pierceable membrane; and iii) a removable membrane provided with means for removing the membrane by pulling it off.

1/1




Fig. 1




Fig. 2

SUBSTITUTE SHEET (RULE 26)

INTERNATIONAL SEARCH REPORT

tional Application No
PCT/NL 99/00239

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 A01N3/02 A47G7/02 B65D85/50 A01G5/06

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 A01N A47G A01G B65D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	DATABASE WPI Section Ch, Week 9045 Derwent Publications Ltd., London, GB; Class A97, AN 90-340094 XP002089962 & JP 02 247101 A (GEM KK), 2 October 1990 (1990-10-02) abstract	1-3, 5, 8
Y		4, 6, 7
Y	DATABASE WPI Section Ch, Week 9717 Derwent Publications Ltd., London, GB; Class G04, AN 97-186861 XP002089963 & JP 09 047154 A (FUSHIMI SEISAKUSHO KK), 18 February 1997 (1997-02-18) abstract	4
	-/-	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubt on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

11 August 1999

Date of mailing of the international search report

19/08/1999

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040. Tx. 31 651 epo nl.
Fax: (+31-70) 340-3016

Authorized officer

Lamers, W

INTERNATIONAL SEARCH REPORT

International Application No
PCT/NL 99/00239

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 2 971 292 A (G.J.MALECKI) 14 February 1961 (1961-02-14) column 1, line 70 - line 72 column 2, line 10 - column 3, line 3 column 6, line 38 - column 7, line 15 column 7, line 32 - line 34 column 9, line 13 - line 16	1-3
Y	---	6
X	US 4 906 276 A (HUGHES JOHN) 6 March 1990 (1990-03-06) column 1, line 6 - line 10 column 3, line 3 - line 11 column 4, line 23 - line 30 column 6, line 38 - line 42 column 11, line 21 - line 24	1-4
Y	---	
Y	US 4 225 679 A (PILATO LOUIS A) 30 September 1980 (1980-09-30) column 1, line 11 - line 15 column 3, line 15	4
X	---	
X	DATABASE WPI Section Ch, Week 8833 Derwent Publications Ltd., London, GB; Class A86, AN 88-230437 XP002089965 & JP 63 162601 A (NIPPON SHOKUBAI KAGAKU KOGYO CO LTD), 6 July 1988 (1988-07-06) abstract	1-3
X	---	
X	GB 2 215 174 A (BARCLAY WILLIAM MATHIESON; STENHOUSE PETER WILLIAM FRANCI) 20 September 1989 (1989-09-20) the whole document	1,7-10
X	---	
X	DATABASE WPI Section PQ, Week 9637 Derwent Publications Ltd., London, GB; Class P13, AN 96-365353 XP002111827 & JP 08 172911 A (NIPPON TURNER KK), 9 July 1996 (1996-07-09) abstract see figure	9,10
Y	---	7
X	WO 98 16428 A (QUIDING DOUGLAS C) 23 April 1998 (1998-04-23) page 5, line 1 - line 5 figure 3	9,10

	-/--	

INTERNATIONAL SEARCH REPORT

International Application No
PCT/NL 99/00239

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>DATABASE WPI Section PQ, Week 9634 Derwent Publications Ltd., London, GB; Class P27, AN 96-334334 XP002089966 & BE 1 008 661 A (SCHORPION E M P), 2 July 1996 (1996-07-02) abstract</p> <p>---</p>	9,10
X	<p>US 5 580 975 A (TADA SUGURU) 3 December 1996 (1996-12-03) column 1, line 6 - line 11 column 1, line 55 - column 2, line 57 column 3, line 9 - column 4, line 6 column 4, line 34 - line 51 figures 3,4</p> <p>---</p>	9
A	<p>DATABASE WPI Section PQ, Week 8607 Derwent Publications Ltd., London, GB; Class P27, AN 86-046737 XP002089964 & SE 8 403 211 A (NILSSON-ENGERTH B), 15 December 1985 (1985-12-15) cited in the application</p> <p>-----</p>	1-10

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/NL 99/00239

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
JP 2247101 A	02-10-1990	NONE	
JP 9047154 A	18-02-1997	NONE	
US 2971292 A		NONE	
US 4906276 A	06-03-1990	US 4985062 A US 4985061 A	15-01-1991 15-01-1991
US 4225679 A	30-09-1980	NONE	
JP 63162601 A	06-07-1988	NONE	
GB 2215174 A	20-09-1989	NONE	
JP 8172911 A	09-07-1996	NONE	
WO 9816428 A	23-04-1998	AU 7515896 A	11-05-1998
BE 1008661 A	02-07-1996	NONE	
US 5580975 A	03-12-1996	JP 2509083 B JP 7082101 A NL 9401194 A	19-06-1996 28-03-1995 16-02-1995
SE 8403211 A	15-12-1985	NONE	

THIS PAGE BLANK (USPTO)

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☒ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☒ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.

THIS PAGE BLANK (USPTO)